



男：46岁 脱发时间：10年以上 使用过米诺地尔

4次资料前拍摄图

8次治疗后前拍摄图

Clinical Case – Wenzhou, China

Male, Age 46

Hair loss duration: Over 10 years

Previously used medication: Minoxidil

Before treatment (photo) / After 8 sessions (photo)

1. Hair loss medications used before treatment:

- Minoxidil
 - Usage duration: 6 months
 - Effect: No noticeable visible effect

2. After 8 sessions of AMURUN treatment:

- Nurses and staff noted significant improvement compared to before.
- More effective than Minoxidil; hopeful for further improvements.



男：40岁 脱发时间：20年以上，之前服用过非那雄胺片

治疗3周后实际效果

治疗6周后实际效果，
8次治疗结束后根据效果再次确定增加次数

Clinical Case – Wenzhou, China

Male, Age 40

Hair loss duration: Over 20 years

Previously used medication: Finasteride (Propecia)

After 3 weeks of treatment / After 6 weeks of treatment

1. Hair loss medications used before treatment:

- Minoxidil (1 year ago), Propecia (2 years ago)
 - Usage duration: Over 2 years
 - Effect: No significant effect despite prolonged use; discontinued.

2. After 8 sessions of AMURUN treatment:

- Continued up to 9 sessions; significant changes in hair condition.
- Future follow-up planned to monitor ongoing changes.



男：31岁 脱发时间：5年以上,之前使用过非那雄安片

治疗5周后实际效果

治疗8周后实际效果,
治疗结束：观察后增加治疗

Clinical Case – Wenzhou, China

Male, Age 31

Hair loss duration: Over 5 years

Previously used medication: Finasteride (Propecia)

After 5 weeks of treatment / After 8 weeks of treatment

1. Hair loss medications used before treatment:

- Propecia
 - Usage duration: Over 1 year
 - Effect: Aware of side effects, but continued due to some effectiveness; also used anti-hair loss shampoo consistently.

2. After 8 sessions of AMURUN treatment:

- Noticed significant improvements.
- According to the manufacturer, effects appear in stages; this has been the most effective and fastest-acting treatment used so far.
- Hopes for even better results going forward.